

KI SOME NOTES ON THE INCLUSION BETWEEN MORREY SPACES

by Nicky Tumalun

Submission date: 03-Jul-2023 01:21PM (UTC+0700)

Submission ID: 2125895307

File name: alun_KI_SOME_NOTES_ON_THE_INCLUSION_BETWEEN_MORREY_SPACES....pdf (141.47K)

Word count: 2876

Character count: 11202

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/359481329>

1 Some notes on the inclusion between Morrey spaces

Article in *Journal of Mathematical Inequalities* · March 2022

DOI:10.7153/jmi-2022-16-26

CITATIONS

2

READS

80

2 authors, including:



Nicky Kurnia Tumulun
Universitas Negeri Manado

10 PUBLICATIONS 23 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



Konsep Sudut pada Ruang Bernorma-n dan Aplikasi Kelas Stummel dalam Teori Regularitas [View project](#)

All content following this page was uploaded by Nicky Kurnia Tumulun on 26 March 2022.

The user has requested enhancement of the downloaded file.

SOME NOTES ON THE INCLUSION BETWEEN MORREY SPACES

PHILOTHEUS E. A. TUERAH AND NICKY K. TUMALUN*

(Communicated by L. Liu)

Abstract. In this paper, we show that the Morrey spaces $\mathcal{M}_{q_1}^p(\mathbb{R}^n)$ cannot be contained in the weak Morrey spaces $w\mathcal{M}_{q_2}^p(\mathbb{R}^n)$ for $q_1 \neq q_2$. We also show that the vanishing Morrey spaces $\mathcal{V}\mathcal{M}_q^p(\mathbb{R}^n)$ are not empty and properly contained in the Morrey spaces $\mathcal{M}_q^p(\mathbb{R}^n)$.

1. Introduction

Let $1 \leq p \leq q < \infty$ and $n \geq 2$. The Morrey space $\mathcal{M}_q^p(\mathbb{R}^n)$ is the set of all functions $f \in L_{\text{loc}}^p(\mathbb{R}^n)$ for which

$$\|f\|_{\mathcal{M}_q^p} = \sup_{x \in \mathbb{R}^n, r > 0} |B(x, r)|^{\frac{1}{q} - \frac{1}{p}} \|f\|_{L^p(B(x, r))} < \infty,$$

where

$$\|f\|_{L^p(B(x, r))} = \left(\int_{B(x, r)} |f(y)|^p dy \right)^{\frac{1}{p}}.$$

Here $B(x, r)$ is the open ball in Euclidean space \mathbb{R}^n with center x and radius r , and $|B(x, r)|$ denotes its Lebesgue measure. Meanwhile, the weak Morrey space $w\mathcal{M}_q^p(\mathbb{R}^n)$ is defined to be the set of all functions $f \in wL_{\text{loc}}^p(\mathbb{R}^n)$ for which

$$\|f\|_{w\mathcal{M}_q^p} = \sup_{x \in \mathbb{R}^n, r > 0} |B(x, r)|^{\frac{1}{q} - \frac{1}{p}} \|f\|_{wL^p(B(x, r))} < \infty,$$

where

$$\|f\|_{wL^p(B(x, r))} = \sup_{t > 0} t |\{y \in B(x, r) : |f(y)| > t\}|^{\frac{1}{p}},$$

and $|\{y \in B(x, r) : |f(y)| > t\}|$ also denotes the Lebesgue measure of the set $\{y \in B(x, r) : |f(y)| > t\}$. Now, we define

$$\mathcal{V}\mathcal{M}_q^p(\mathbb{R}^n) = \left\{ f \in \mathcal{M}_q^p(\mathbb{R}^n) : \lim_{r \rightarrow 0} \mathcal{M}_f(r) = 0 \right\},$$

Mathematics subject classification (2020): 46E30, 42B35.

Keywords and phrases: Morrey spaces, vanishing Morrey spaces.

* Corresponding author.

where

$$\mathcal{M}_f(r) = \sup_{x \in \mathbb{R}^n} |B(x, r)|^{\frac{1}{q} - \frac{1}{p}} \|f\|_{L^p(B(x, r))}.$$

The set $\mathcal{V}\mathcal{M}_q^p(\mathbb{R}^n)$ is called the *vanishing Morrey space*. It is clear that $\mathcal{V}\mathcal{M}_q^p(\mathbb{R}^n)$ is a subset of $\mathcal{M}_q^p(\mathbb{R}^n)$.

The Morrey spaces were introduced by C. B. Morrey [1] and the vanishing Morrey spaces were introduced in [2]. Recently, many authors are attracted in studying the inclusion properties between Morrey spaces [3, 4, 5, 6, 7, 8]. One interesting result stated in [5, Remark 4.5], that is, the weak Morrey spaces $w\mathcal{M}_{q_1}^p(\mathbb{R}^n)$ cannot be contained in the weak Morrey spaces $w\mathcal{M}_{q_2}^p(\mathbb{R}^n)$ and vice versa, for distinct values q_1 and q_2 . This statement was deduced by a characterization of inclusion between weak Morrey spaces and its parameters, which is proved by using Closed Graph Theorem and Morrey norm estimate for the characteristic functions of balls [5, Theorem 4.4]. Regarding to the inclusion between vanishing Morrey spaces and Morrey spaces over a bounded domain, it was stated in [9] that the vanishing Morrey spaces are properly contained in the Morrey spaces without giving an explicit counter example.

In this paper, we will prove that the Morrey spaces $\mathcal{M}_{q_1}^p(\mathbb{R}^n)$ cannot be contained in the weak Morrey spaces $w\mathcal{M}_{q_2}^p(\mathbb{R}^n)$ and the Morrey spaces $\mathcal{M}_{q_2}^p(\mathbb{R}^n)$ cannot be contained in the weak Morrey spaces $w\mathcal{M}_{q_1}^p(\mathbb{R}^n)$, for different values q_1 and q_2 . This result is more general and sharp than the previous result in [5] since we can recover that previous result and the fact that $\mathcal{M}_q^p(\mathbb{R}^n)$ is a proper subset of $w\mathcal{M}_q^p(\mathbb{R}^n)$ [6, Theorem 1.2]. We also note that our method here is different than in [5] because we give a function which belongs to $\mathcal{M}_{q_1}^p(\mathbb{R}^n) \setminus w\mathcal{M}_{q_2}^p(\mathbb{R}^n)$ and a function which belongs to $\mathcal{M}_{q_2}^p(\mathbb{R}^n) \setminus w\mathcal{M}_{q_1}^p(\mathbb{R}^n)$. Furthermore, by using the idea in [8], we also show that the vanishing Morrey spaces $\mathcal{V}\mathcal{M}_q^p(\mathbb{R}^n)$ are non empty and properly contained in $\mathcal{M}_q^p(\mathbb{R}^n)$ by providing some examples.

The positive constant C that appears in the proofs of all theorems may vary from line to line and the notation $C = C(n, p, q)$ indicates that C depends only on n, p and q .

2. A note on the inclusion between weak Morrey spaces

Let $1 \leq p < q < \infty$ and $\gamma = \frac{n}{q} < n$. Define a function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ by the formula

$$f(y) = \begin{cases} |y|^{-\gamma}, & y \neq 0, \\ 0, & y = 0. \end{cases} \quad (1)$$

It is clear that $1 - \frac{n}{p\gamma} < 0$ and $n - p\gamma > 0$ by observing to the given assumptions.

The function f , that appears in Lemma 1 and 3, is defined by (1).

LEMMA 1. *If $x \in \mathbb{R}^n$ and $r > 0$, then $\|f\|_{L^p(B(x, r))} \leq Cr^{\frac{n}{p} - \gamma} = Cr^{\frac{n}{p} - \frac{n}{q}}$, where $C = C(n, p, q)$.*

Proof. Note that

$$\begin{aligned} \int_{B(x,r)} |f(y)|^p dy &= \int_{\{|y| \leq |x-y| < r\}} |y|^{-p\gamma} dy + \int_{\{|x-y| < |y|\} \cap \{|x-y| < r\}} |y|^{-p\gamma} dy \\ &= I + II. \end{aligned}$$

Since $n - p\gamma > 0$, we have

$$I \leq \int_{\{|y| < r\}} |y|^{-p\gamma} dy = C \int_0^r t^{n-p\gamma-1} dt = Cr^{n-p\gamma}$$

and

$$II \leq \int_{\{|x-y| < r\}} |x-y|^{-p\gamma} dy = C \int_0^r t^{n-p\gamma-1} dt = Cr^{n-p\gamma},$$

by using polar coordinate for radial function. Therefore

$$\|f\|_{L^p(B(x,r))} = \left(\int_{B(x,r)} |f(y)|^p dy \right)^{\frac{1}{p}} \leq C \left(Cr^{\frac{n}{p}-\gamma} \right)^{\frac{1}{p}} = Cr^{\frac{n}{p}-\frac{n}{q}},$$

which proves the lemma. \square

The following lemma is not hard to prove. We leave its proof to the reader.

LEMMA 2. Let $s > 0$, $M \geq 0$, and $\varphi : (0, \infty) \rightarrow [0, \infty)$. If

$$\sup_{0 < t \leq s} \varphi(t) = M = \sup_{s < t < \infty} \varphi(t),$$

then

$$\sup_{t > 0} \varphi(t) = M.$$

Using the above lemma, we can compute the weak Lebesgue norm of f on the ball $B(0, r)$ with arbitrary radius r .

LEMMA 3. If $r > 0$, then $\|f\|_{wL^p(B(0,r))} = Cr^{-\gamma+\frac{n}{p}} = Cr^{-\frac{n}{q}+\frac{n}{p}}$, where $C = C(n, p, q)$.

Proof. Let r be an arbitrary positive real number. Note that, for every $t > 0$, we have

$$\begin{aligned} |\{y \in B(0, r) : |f(y)| > t\}| &= \left| \left\{ y \in B(0, r) : |y| < t^{-\frac{1}{\gamma}} \right\} \right| \\ &= \left| B(0, r) \cap B(0, t^{-\frac{1}{\gamma}}) \right|. \end{aligned} \quad (2)$$

We now define $\varphi : (0, \infty) \rightarrow [0, \infty)$ by the formula

$$\varphi(t) = t |\{y \in B(0, r) : |f(y)| > t\}|^{\frac{1}{p}}. \quad (3)$$

For every $t > r^{-\gamma}$, we obtain $t^{-\frac{1}{\gamma}} < r$. Then

$$|\{y \in B(0, r) : |f(y)| > t\}| = |B(0, t^{-\frac{1}{\gamma}})| = Ct^{-\frac{n}{\gamma}},$$

by using (2). This gives us

$$t|\{y \in B(0, r) : |f(y)| > t\}|^{\frac{1}{p}} = Ct \left(t^{-\frac{n}{\gamma}}\right)^{\frac{1}{p}} = Ct^{1-\frac{n}{p\gamma}}, \quad \forall t \in (r^{-\gamma}, \infty). \quad (4)$$

On the other hand, for every $t \leq r^{-\gamma}$, we have $t^{-\frac{1}{\gamma}} \geq r$. Hence

$$|\{y \in B(0, r) : |f(y)| > t\}| = |B(0, r)| = Cr^n,$$

which comes from (2). Therefore,

$$t|\{y \in B(0, r) : |f(y)| > t\}|^{\frac{1}{p}} = Ct(r^n)^{\frac{1}{p}} = Ctr^{\frac{n}{p}}, \quad \forall t \in (0, r^{-\gamma}]. \quad (5)$$

We obtain

$$\varphi(t) = \begin{cases} Ct^{1-\frac{n}{p\gamma}}, & \forall t \in (r^{-\gamma}, \infty) \\ Ctr^{\frac{n}{p}}, & \forall t \in (0, r^{-\gamma}]. \end{cases}$$

by virtue of (4) and (5). Observing φ non increasing on $(r^{-\gamma}, \infty)$ and non decreasing on $(0, r^{-\gamma}]$, since $1 - \frac{n}{p\gamma} < 0$ and $\frac{n}{p} > 0$ respectively, then

$$\sup_{r^{-\gamma} < t < \infty} \varphi(t) = Cr^{-\gamma+\frac{n}{p}} = \sup_{0 < t \leq r^{-\gamma}} \varphi(t). \quad (6)$$

Thus

$$\|f\|_{wL^p(B(0,r))} = \sup_{t>0} \varphi(t) = Cr^{-\gamma+\frac{n}{p}} = Cr^{-\frac{n}{q}+\frac{n}{p}},$$

that is concluded from Lemma 2. \square

By taking Lemma 2 and Lemma 3 as the tools, we are ready to state and prove the first main result of this paper.

THEOREM 1. *Let $1 \leq p < q_1 < \infty$ and $1 \leq p < q_2 < \infty$. If $q_1 \neq q_2$, then $\mathcal{M}_{q_1}^p(\mathbb{R}^n) \not\subseteq w\mathcal{M}_{q_2}^p(\mathbb{R}^n)$ and $\mathcal{M}_{q_2}^p(\mathbb{R}^n) \not\subseteq w\mathcal{M}_{q_1}^p(\mathbb{R}^n)$.*

Proof. We will only prove that $\mathcal{M}_{q_1}^p(\mathbb{R}^n)$ is not contained by $w\mathcal{M}_{q_2}^p(\mathbb{R}^n)$. The proof that $\mathcal{M}_{q_2}^p(\mathbb{R}^n)$ is not contained by $w\mathcal{M}_{q_1}^p(\mathbb{R}^n)$ can be done by similar method.

Let $\gamma_1 = n/q_1$ and $f_1 : \mathbb{R}^n \rightarrow \mathbb{R}$, defined by the formula

$$f_1(y) = \begin{cases} |y|^{-\gamma_1}, & y \neq 0, \\ 0, & y = 0. \end{cases}$$

We will show that $f_1 \in \mathcal{M}_{q_1}^p(\mathbb{R}^n) \setminus w\mathcal{M}_{q_2}^p(\mathbb{R}^n)$. Let $x \in \mathbb{R}^n$ and $r > 0$ be arbitrarily given. According to Lemma 1, by replacing γ with γ_1 , we obtain

$$|B(x, r)|^{\frac{1}{q_1} - \frac{1}{p}} \|f_1\|_{L^p(B(x,r))} \leq Cr^{\frac{n}{q_1} - \frac{n}{p}} r^{\frac{n}{p} - \frac{n}{q_1}} = C < \infty.$$

This gives us

$$\|f_1\|_{\mathcal{M}_{q_1}^p} = \sup_{x \in \mathbb{R}^n, r > 0} |B(x, r)|^{\frac{1}{q_1} - \frac{1}{p}} \|f_1\|_{L^p(B(x, r))} < \infty,$$

since x and r are arbitrary. Whence $f_1 \in \mathcal{M}_{q_1}^p(\mathbb{R}^n)$. By virtue to Lemma 3, we have

$$\|f_1\|_{w.\mathcal{M}_{q_2}^p} \geq |B(0, r)|^{\frac{1}{q_2} - \frac{1}{p}} \|f_1\|_{wL^p(B(0, r))} = Cr^{\frac{n}{q_2} - \frac{n}{p} - \frac{n}{q_1}} = Cr^n \left(\frac{1}{q_2} - \frac{1}{q_1} \right).$$

Hence $\|f_1\|_{w.\mathcal{M}_{q_2}^p} = \infty$. This is due to arbitrary r and $q_1 \neq q_2$. We conclude that $f_1 \notin w.\mathcal{M}_{q_2}^p(\mathbb{R}^n)$. Thus, we have already proved that $\mathcal{M}_{q_1}^p(\mathbb{R}^n) \not\subseteq w.\mathcal{M}_{q_2}^p(\mathbb{R}^n)$. \square

As an immediate consequence of Theorem 1, we recover the result from [5] which is stated in the following corollary.

COROLLARY 1. *Let $1 \leq p < q_1 < \infty$ and $1 \leq p < q_2 < \infty$. If $q_1 \neq q_2$, then $w.\mathcal{M}_{q_1}^p(\mathbb{R}^n) \not\subseteq w.\mathcal{M}_{q_2}^p(\mathbb{R}^n)$ and $w.\mathcal{M}_{q_2}^p(\mathbb{R}^n) \not\subseteq w.\mathcal{M}_{q_1}^p(\mathbb{R}^n)$.*

3. A note on the inclusion between Morrey spaces and vanishing Morrey spaces

Let $1 \leq p < q < \infty$ and $\delta = \exp(\frac{-2q}{np})$. Define a function $g : \mathbb{R}^n \rightarrow \mathbb{R}$ by the formula

$$g(y) = \begin{cases} \left(\frac{\chi_B(y)}{|y|^{\frac{np}{q}} (\ln|y|)^2} \right)^{\frac{1}{p}}, & y \neq 0, \\ 0, & y = 0, \end{cases} \quad (7)$$

where χ_B is a characteristic function defined on $B = B(0, \delta)$.

The function g in the following lemma is defined by (7). This following lemma shows that the vanishing Morrey spaces in a non empty set.

LEMMA 4. $g \in \mathcal{V}\mathcal{M}_q^p(\mathbb{R}^n)$.

Proof. Let $x \in \mathbb{R}^n$ and $r > 0$ be arbitrarily given. Note that

$$\begin{aligned} |B(x, r)|^{\frac{1}{q} - \frac{1}{p}} \|g\|_{L^p(B(x, r))} &\leq C \left(\int_{|y| \leq |x-y| < r} \frac{\chi_B(y)}{|x-y|^{n-\frac{np}{q}} |y|^{\frac{np}{q}} (\ln|y|)^2} dy \right)^{\frac{1}{p}} \\ &\quad + C \left(\int_{\substack{\{|x-y| < |y|\} \\ \cap \{|x-y| < r\}}} \frac{\chi_B(y)}{|x-y|^{n-\frac{np}{q}} |y|^{\frac{np}{q}} (\ln|y|)^2} dy \right)^{\frac{1}{p}} \\ &= I + II. \end{aligned} \quad (8)$$

Now we have two cases, that is, $\delta \leq r$ or $r < \delta$. Assume $\delta \leq r$, then we have

$$I \leq \int_{|y| < r} \frac{\chi_B(y)}{|y|^n (\ln|y|)^2} dy = \int_{|y| < \delta} \frac{1}{|y|^n (\ln|y|)^2} dy = C \left(\frac{-1}{\ln(\delta)} \right), \quad (9)$$

and

$$\begin{aligned} H &= \int_{\substack{\{|x-y|<|y|<\delta\} \\ \cap \{|x-y|<r\}}} \frac{1}{|x-y|^{n-\frac{np}{q}} |y|^{\frac{np}{q}} (\ln|y|)^2} dy \leq \int_{|x-y|<\delta} \frac{1}{|x-y|^n (\ln|x-y|)^2} dy \\ &= C \left(\frac{-1}{\ln(\delta)} \right), \end{aligned} \quad (10)$$

since $1/t^{np/q}(\ln(t))^2$ decreasing on interval $(0, \delta)$. Assume $r < \delta$. We have

$$I \leq \int_{|y|<r} \frac{1}{|y|^n (\ln|y|)^2} dy = C \left(\frac{-1}{\ln(r)} \right) \leq C \left(\frac{-1}{\ln(\delta)} \right), \quad (11)$$

and

$$\begin{aligned} H &= \int_{\substack{\{|x-y|<|y|<r\} \\ \cap \{|x-y|<r\}}} \frac{1}{|x-y|^{n-\frac{np}{q}} |y|^{\frac{np}{q}} (\ln|y|)^2} dy \leq \int_{|x-y|<r} \frac{1}{|x-y|^n (\ln|x-y|)^2} dy \\ &= C \left(\frac{-1}{\ln(r)} \right) \leq C \left(\frac{-1}{\ln(\delta)} \right), \end{aligned} \quad (12)$$

since $1/t^{np/q}(\ln(t))^2$ decreasing on interval $(0, r) \subseteq (0, \delta)$. By virtue of (8), (9), (10), (11), and (12), we conclude that

$$|B(x, r)|^{\frac{1}{q}-\frac{1}{p}} \|g\|_{L^p(B(x, r))} \leq I + H \leq C \left(\frac{-1}{\ln(\delta)} \right)^{\frac{1}{p}},$$

where $C = C(n, p, q)$. This means $g \in \mathcal{M}_q^p(\mathbb{R}^n)$. We remaind to prove

$$\lim_{r \rightarrow 0} \mathcal{M}_f(r) = 0. \quad (13)$$

For every $0 < r < \delta$, we have shown that

$$\mathcal{M}_f(r) \leq C \left(\frac{-1}{\ln(r)} \right)^{\frac{1}{p}}.$$

This means (13) holds and the proof is done. \square

Now we define a function that will play as an element of Morrey spaces but not in the vanishing Morrey spaces. Let $1 \leq p < q < \infty$. For every $k \in \mathbb{N}$, with $k \geq 3$, we set $x_k = (2^{-k}, \dots, 0) \in \mathbb{R}^n$ and

$$u_k(y) = \begin{cases} 8^{\frac{np}{q}k}, & y \in B(x_k, 8^{-k}), \\ 0, & y \notin B(x_k, 8^{-k}). \end{cases}$$

Define a function $u : \mathbb{R}^n \rightarrow \mathbb{R}$ by the formula

$$u(y) = \left(\sum_{k=3}^{\infty} u_k(y) \right)^{\frac{1}{p}}. \quad (14)$$

We first claim that u belongs to the Morrey spaces $\mathcal{M}_q^p(\mathbb{R}^n)$.

LEMMA 5. $u \in \mathcal{M}_q^p(\mathbb{R}^n)$.

Proof. Let $x \in \mathbb{R}^n$ and $r > 0$ be arbitrarily given. There are two cases: (i) $x \notin B(x_k, 2(4^{-k}))$ for every $k \geq 3$, or, (ii) $x \in B(x_j, 2(4^{-j}))$ for some $j \geq 3$. Assume (i) holds. Then

$$2(4^{-k}) \leq |x - x_k| \leq |x - y| + |y - x_k| < r + 4^{-k},$$

for every $y \in B(x, r) \cap B(x_k, 8^{-k})$. This means $r^{\frac{np}{q}-n} \leq 4^{(n-\frac{np}{q})k}$ and

$$\begin{aligned} r^{\frac{np}{q}-n} \int_{B(x,r)} |u(y)|^p dy &\leq \sum_{k=3}^{\infty} 4^{(n-\frac{np}{q})k} \int_{B(x,r) \cap B(x_k, 8^{-k})} 8^{\frac{np}{q}k} dy \\ &\leq C \sum_{k=3}^{\infty} 2^{(\frac{np}{q}-n)k} < \infty, \end{aligned} \quad (15)$$

where C depends on n . Assume (ii) holds. Since $\{B(x_k, 2(4^{-k}))\}_{k \geq 3}$ is a disjoint collection, then there is only one $j \geq 3$ such that $x \in B(x_j, 2(4^{-j}))$ and $x \notin B(x_k, 2(4^{-k}))$ for every $k \geq 3$ with $k \neq j$. Note that

$$r^{\frac{np}{q}-n} \int_{B(x,r) \cap B(x_j, 8^{-j})} u_j(y) dy = r^{\frac{np}{q}-n} \int_{B(x,r) \cap B(x_j, 8^{-j})} 8^{\frac{np}{q}j} dy \leq C < \infty, \quad (16)$$

where C depends on n, p , and q . By virtue of (16) and the computation of (15), we have

$$\begin{aligned} r^{\frac{np}{q}-n} \int_{B(x,r)} |u(y)|^p dy &= r^{\frac{np}{q}-n} \sum_{k=3}^{\infty} \int_{B(x,r) \cap B(x_k, 8^{-k})} u_k(y) dy \\ &= r^{\frac{np}{q}-n} \int_{B(x,r) \cap B(x_j, 8^{-j})} u_j(y) dy \\ &\quad + r^{\frac{np}{q}-n} \sum_{\substack{k=3 \\ k \neq j}}^{\infty} \int_{B(x,r) \cap B(x_k, 8^{-k})} u_k(y) dy \\ &\leq C + C \sum_{\substack{k=3 \\ k \neq j}}^{\infty} 2^{(\frac{np}{q}-n)k} < \infty, \end{aligned} \quad (17)$$

where C depends on n, p , and q . Combining (16) and (17), whence

$$|B(x, r)|^{\frac{1}{q}-\frac{1}{p}} \|u\|_{L^p(B(x,r))} = C \left(r^{\frac{np}{q}-n} \int_{B(x,r)} |u(y)|^p dy \right)^{\frac{1}{p}} \leq C < \infty,$$

where C depends on n, p , and q . Therefore $u \in \mathcal{M}_q^p(\mathbb{R}^n)$. \square

The following theorem states that the vanishing Morrey spaces is a non empty proper subset of the Morrey spaces. This theorem is the second main result in this paper.

THEOREM 2. *Let $1 \leq p < q < \infty$. Then $\mathcal{V}\mathcal{M}_q^p(\mathbb{R}^n)$ is a non empty proper subset of $\mathcal{M}_q^p(\mathbb{R}^n)$.*

Proof. According to Lemma 4, $\mathcal{V}\mathcal{M}_q^p(\mathbb{R}^n)$ is non empty, and according to Lemma 5, the function u belongs to $\mathcal{M}_q^p(\mathbb{R}^n)$. Therefore, we need only to show that u does not belong to $\mathcal{V}\mathcal{M}_q^p(\mathbb{R}^n)$. Let $0 < r < 1$. By the Archimedean property, there is an integer $k \geq 3$ such that $8^{-k} < r$. Then

$$\begin{aligned} (\mathcal{M}_f(r))^p &\geq Cr^{\frac{np}{q}-n} \int_{B(x_k, r)} |u(y)|^p dy \geq C \int_{B(x_k, 8^{-k})} u_k(y) dy \\ &= C \int_{B(x_k, 8^{-k})} 8^{\frac{np}{q}k} dy \geq C8^{-nk} \int_{B(x_k, 8^{-k})} 1 dy = C > 0, \end{aligned}$$

where C depends on n . This means $\mathcal{M}_f(r)$ is bounded away from zero as r tends to zero. Thus $u \notin \mathcal{V}\mathcal{M}_q^p(\mathbb{R}^n)$. \square

REFERENCES

- [1] C. B. MORREY, *On the solutions of quasi-linear elliptic partial differential equations*, Trans. Amer. Math. Soc., **43**, (1938), 126–166.
- [2] C. VITANZA, *Functions with vanishing Morrey norm and elliptic partial differential equations*, in Proceedings of Methods of Real Analysis and Partial Differential Equations, Capri, (1990).
- [3] D. I. HAKIM AND Y. SAWANO, *Complex interpolation of various subspaces of Morrey spaces*, Sci. China Math., **63**, 5 (2020), 937–964.
- [4] H. GUNAWAN, D. I. HAKIM, AND M. IDRIS, *Proper inclusions of Morrey spaces*, Glasnik Matematički, **53**, 1 (2018), 143–151.
- [5] H. GUNAWAN, D. I. HAKIM, K. M. LIMANTA, AND A. A. MASTA, *Inclusion properties of generalized Morrey spaces*, Math. Nachr., **290**, 1 (2017), 332–340.
- [6] H. GUNAWAN, D. I. HAKIM, E. NAKAI, AND Y. SAWANO, *On inclusion relation between weak Morrey spaces and Morrey spaces*, Nonlinear Analysis., **168**, (2018), 27–31.
- [7] N. K. TUMALUN AND H. GUNAWAN, *Morrey spaces are embedded between weak Morrey spaces and Stummel classes*, J. Indones. Math. Soc., **253**, (2019), 203–209.
- [8] N. K. TUMALUN, D. I. HAKIM, AND H. GUNAWAN, *Inclusion between generalized Stummel classes and other function spaces*, Math. Inequal. Appl., **232**, (2020), 547–562.
- [9] N. SAMKO, *Maximal, potential and singular operators in vanishing generalized Morrey spaces*, J. Global. Optim., **57**, (2013), 1385–1399.

(Received March 22, 2021)

Philotheus E. A. Tuerah
Mathematics Department, Faculty of Mathematics and Natural Sciences
Universitas Negeri Manado, Tondano Selatan District, Minahasa Regency
Sulawesi Utara Province, Indonesia, Indonesia
e-mail: pheaturah@unima.ac.id

Nicky K. Tumulun
Mathematics Department, Faculty of Mathematics and Natural Sciences
Universitas Negeri Manado, Tondano Selatan District, Minahasa Regency
Sulawesi Utara Province, Indonesia
e-mail: nickyktumulun@unima.ac.id

KI SOME NOTES ON THE INCLUSION BETWEEN MORREY SPACES

ORIGINALITY REPORT

21 %
SIMILARITY INDEX

21 %
INTERNET SOURCES

0 %
PUBLICATIONS

0 %
STUDENT PAPERS

PRIMARY SOURCES

1 ojs3.unpatti.ac.id
Internet Source

21 %

Exclude quotes Off

Exclude matches < 2%

Exclude bibliography On